



Cambridge International Examinations
Cambridge Ordinary Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER



CHEMISTRY

5070/31

Paper 3 Practical Test

October/November 2016

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

Qualitative Analysis Notes are printed on page 8.

You should show the essential steps in any calculations and record experimental results in the spaces provided on the Question Paper.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
Total	

This document consists of **6** printed pages and **2** blank pages.

- 1 Mangalloy is made by alloying steel with manganese. The percentage by mass of manganese in mangalloy can be determined by converting all the manganese in the alloy into aqueous manganate(VII) ions and then using the solution to titrate aqueous iron(II) sulfate.

No indicator is needed for this titration as the products of the reaction are almost colourless and one drop of aqueous manganate(VII) ions in excess produces a permanent pale pink colour.

P is an aqueous solution of manganate(VII) ions, MnO_4^- . The solution was prepared by converting all the manganese in a 2.12 g sample of mangalloy into manganate(VII) ions and making the final volume up to 250 cm^3 by adding water.

Q is 0.0800 mol/dm^3 iron(II) sulfate.

- (a) Put **P** into the burette.

The colour of **P** makes it difficult to see the bottom of the meniscus so you should take all your readings using the top of the meniscus.

Pipette a 25.0 cm^3 (or 20.0 cm^3) portion of **Q** into a flask.

Add **P** from the burette. At first the purple colour disappears quickly but as more **P** is added the colour disappears less quickly. At the end-point, one drop of **P** produces a pale pink colour that does not disappear on swirling.

Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

Results

Burette readings

titration number	1	2	
final reading / cm^3			
initial reading / cm^3			
volume of P used / cm^3			
best titration results (✓)			

Summary

Tick (✓) the best titration results.

Using these results, the average volume of **P** required was cm^3 .

Volume of **Q** used was cm^3 .

[12]

- (b) **Q** is 0.0800 mol/dm^3 iron(II) sulfate.

Calculate the number of moles of iron(II) sulfate present in the volume of **Q** used.

number of moles of iron(II) sulfate[1]

- (c) Using your answer from (b), calculate the number of moles of manganate(VII) ions present in the average volume of **P** required.

[Five moles of iron(II) sulfate react with one mole of manganate(VII) ions.]

number of moles of manganate(VII) ions[1]

- (d) Using your answer from (c), calculate the number of moles of manganate(VII) ions in 250 cm^3 of **P**.

number of moles of manganate(VII) ions in 250 cm^3 of **P**[1]

- (e) Using your answer from (d), calculate the mass of manganese in the 2.12g sample of mangalloy.

[The relative atomic mass of manganese is 55.]

mass of manganese in 2.12g of mangalloy g [1]

- (f) Using your answer from (e), calculate the percentage by mass of manganese in mangalloy.

percentage by mass of manganese in mangalloy % [1]

[Total: 17]

- 2 You are provided with solution **R** and solid **S**.
Carry out the following tests and record your observations in the table.
You should test and name any gas evolved.

test no.	test	observations
1	<p>(a) To 1 cm depth of R in a boiling tube, add 2 drops of litmus solution.</p> <p>(b) To the mixture from (a), add about twice the volume of aqueous sodium hydroxide.</p> <p>Keep this mixture for use in test 2.</p>	
2	To the mixture from test 1, add a piece of aluminium and warm the boiling tube until reaction just begins.	
3	<p>(a) To 1 cm depth of R in a boiling tube, add an equal volume of aqueous iron(II) sulfate. Heat the mixture until the liquid just begins to boil.</p> <p>(b) To the mixture from (a), add aqueous sodium hydroxide until no further change occurs.</p>	
4	To 2 cm depth of R in a test-tube, add a piece of magnesium.	

test no.	test	observations
5	To 2 cm depth of R in a test-tube, add a small amount of S . Keep this solution for use in tests 6 and 7 .	
6	Transfer about half of the solution from test 5 to a test-tube and add aqueous sodium hydroxide until no further change occurs.	
7	To the rest of the solution from test 5 in a test-tube, add aqueous ammonia until no further change occurs.	
8	To a small amount of S in a hard-glass test-tube, add an equal amount of ammonium chloride and mix the solids together. Heat the mixture.	

[19]

Conclusions

Give the formula of a cation and the formula of an anion in **R**.

The formula of a cation in **R** is

The formula of an anion in **R** is

Give the formula of a cation and the formula of an anion in **S**.

The formula of a cation in **S** is

The formula of an anion in **S** is

[4]

[Total: 23]

QUALITATIVE ANALYSIS NOTES

Tests for anions

<i>anion</i>	<i>test</i>	<i>test result</i>
carbonate (CO_3^{2-})	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl^-) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I^-) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO_3^-) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO_4^{2-}) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.

Tests for aqueous cations

<i>cation</i>	<i>effect of aqueous sodium hydroxide</i>	<i>effect of aqueous ammonia</i>
aluminium (Al^{3+})	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH_4^+)	ammonia produced on warming	–
calcium (Ca^{2+})	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper(II) (Cu^{2+})	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe^{2+})	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe^{3+})	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn^{2+})	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia (NH_3)	turns damp red litmus paper blue
carbon dioxide (CO_2)	turns limewater milky
chlorine (Cl_2)	bleaches damp litmus paper
hydrogen (H_2)	'pops' with a lighted splint
oxygen (O_2)	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.